

Linear Algebra

[KOMS120301] - 2023/2024

3.1 - Linear System of Equations

Dewi Sintiar

Computer Science Study Program
Universitas Pendidikan Ganesha

Week 3 (September 2023)

Motivating example



Rp 21.000,00



Rp 22.000,00



???

Motivating example



Rp 26.000,00



Rp 24.500,00



Rp 16.000,00



???

Part 1: System of linear equations

(We sometime call it "linear system")

Learning objectives

After this lecture, you should be able to:

1. analyze the components of a system of linear equations;
2. verify whether a given set is a solution of a linear system;
3. identify a homogeneous and non-homogeneous linear system;
4. formulate the coefficient matrix and augmented matrix of a given linear system;
5. showing that elementary row system gives an equivalent linear system;
6. analyze the geometric interpretation of a linear system with 1, 2, or 3 variables;
7. apply the elimination and substitution algorithms to solve a linear system;
8. explain the concept of linear system written in triangular matrix or in echelon form.

Terminology and notation (1)

Given unknowns variables x_1, x_2, \dots, x_n , a **linear equation** on the variables is defined as:

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = b \quad (1)$$

where $a_1, a_2, \dots, a_n, b \in \mathbb{R}$ (this can be replaced by another *field*).

A **solution** of equation (1) is a **list of values for the unknowns**, or a **vector u in \mathbb{R}^n** .

$$x_1 = r_1, x_2 = r_2, \dots, x_n = r_n \quad \text{or} \quad u = (r_1, r_2, \dots, r_n)$$

This means that the following is correct:

$$a_1r_1 + a_2r_2 + \dots + a_nr_n = b$$

In this case, we say that **u satisfies** equation (1).

Terminology and notation (2)

In equation (1):

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$$

We say that:

- the equation is written in the **standard form**
- the constant a_k is the **coefficient** of x_k
- b is the **constant term** of the equation

Note: If n is small, we use different letters to denote the variables, instead of using indexing.

Example: how many solutions are there?

Given an equation:

$$2x + 3y - z = 4$$

Can you find a solution for the equation?

How many solutions that you can find?

System of linear equations

A **system of linear equations** is a list of linear equations:
 L_1, L_2, \dots, L_m with the same variables x_1, x_2, \dots, x_n .

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \quad (1)$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \quad (2)$$

$$\dots \dots \dots \quad (3)$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \quad (4)$$

where a_{ij} and b_i are constants.

- The system of linear equations is written in **standard form**
- The system is called an **$m \times n$ system**
- a_{ij} is the **coefficient** of variable x_j in the equation L_i
- the number b_i is the **constant** of the equation L_i

What does the word “**linear**” mean???

Linear means



Solution of “system of linear equations”

$$a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2$$

.....

$$a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m$$

A **solution** of the system is a **list of values for the unknowns** or a vector u in \mathbb{R}^n .

Example: verifying solution of a linear system

Given the following system of linear equations:

$$\begin{cases} x_1 + x_2 + 4x_3 + 3x_4 = 5 \\ 2x_1 + 3x_2 + x_3 - 2x_4 = 1 \\ x_1 + 2x_2 - 5x_3 + 4x_4 = 3 \end{cases}$$

- What is the value of m and n in the system?
- Determine whether the following are solutions of the system!
 1. $u = (-8, 6, 1, 1)$
 2. $v = (-10, 5, 1, 2)$

Part 2: Types of system of linear equations

Augmented and coefficient matrices of a system

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & & & \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \cdots & & & & \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{bmatrix}$$

- the left matrix is called the **coefficient matrix** of the system;
- the right matrix is called the **augmented matrix** of the system.

Furthermore, the vector

$$\begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

is called the **constant vector** (or **constant matrix**) of the system.

Example: augmented matrix and coefficient matrix

Given the following system of equations:

$$\begin{cases} x_1 + x_2 + 4x_3 + 3x_4 = 5 \\ 2x_1 + 3x_2 + x_3 - 2x_4 = 1 \\ x_1 + 2x_2 - 5x_3 + 4x_4 = 3 \end{cases}$$

Example: augmented matrix and coefficient matrix

Given the following system of equations:

$$\begin{cases} x_1 + x_2 + 4x_3 + 3x_4 = 5 \\ 2x_1 + 3x_2 + x_3 - 2x_4 = 1 \\ x_1 + 2x_2 - 5x_3 + 4x_4 = 3 \end{cases}$$

The coefficient matrix and the augmented matrix are as follows:

$$\begin{bmatrix} 1 & 1 & 4 & 3 \\ 2 & 3 & 1 & -2 \\ 1 & 2 & -5 & 4 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 1 & 4 & 3 & 5 \\ 2 & 3 & 1 & -2 & 1 \\ 1 & 2 & -5 & 4 & 3 \end{bmatrix}$$

Homogeneous & non-homogeneous linear system

For the given system:

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \cdots & & & & \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & & & \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

It is called **homogeneous** if $b_i = 0, \forall i$. Otherwise, it is called **non-homogeneous**.

Every homogeneous linear system always has a solution. Can you guess what it is?

Degenerate and non-degenerate linear equations

A linear equation is **degenerate** if all coefficients are zero

$$0x_1 + 0x_2 + \cdots + 0x_n = b$$

Can you guess, **what is the condition s.t. the linear equation has a solution?**

Degenerate and non-degenerate linear equations

A linear equation is **degenerate** if all coefficients are zero

$$0x_1 + 0x_2 + \cdots + 0x_n = b$$

Can you guess, **what is the condition s.t. the linear equation has a solution?**

- If $b \neq 0$, then the equation has no solution.
- If $b = 0$, then every vector $u = (r_1, r_2, \dots, r_n)$ in \mathbb{R}^n is a solution.

Degenerate linear equations

Theorem

Let \mathcal{L} be a system of linear equations that contains a degenerate equation L , with constant b .

- 1. If $b \neq 0$, then the system \mathcal{L} has no solution.*
- 2. If $b = 0$, then L may be deleted from \mathcal{L} without changing the solution set of \mathcal{L} .*

Leading unknown in a nondegenerate linear equation

Given a **non-degenerate** linear equation L .

- What can you say about *the coefficients of L* ?

Leading unknown in a nondegenerate linear equation

Given a **non-degenerate** linear equation L .

- What can you say about *the coefficients of L* ?

L has at least one *non-zero coefficient*

Example

The following are non-degenerate linear equations.

$$0x_1 + 0x_2 + 5x_3 + 6x_4 + 0x_5 + 8x_6 = 7 \quad \text{and} \quad 0x + 2y - 4z = 5$$

The zero coefficients are usually omitted.

$$5x_3 + 6x_4 + 8x_6 = 7 \quad \text{and} \quad 2y - 4z = 5$$

Part 3: Elementary row operations

Linear combination

Given:

$$a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \quad (1)$$

$$a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \quad (2)$$

$$\dots\dots\dots \quad (3)$$

$$a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \quad (4)$$

Multiply the m equations by constants c_1, c_2, \dots, c_m :

$$(c_1 a_{11} + \cdots + c_m a_{m1})x_1 + \cdots + (c_1 a_{1n} + \cdots + c_m a_{mn})x_n = c_1 b_1 + \cdots + c_m b_m$$

This is a **linear combination** of the equations in the system.

Example

Given a linear system:

$$\begin{cases} x_1 + x_2 + 4x_3 + 3x_4 = 5 \\ 2x_1 + 3x_2 + x_3 - 2x_4 = 1 \\ x_1 + 2x_2 - 5x_3 + 4x_4 = 3 \end{cases}$$

Then:

$$3L_1 : 3x_1 + 3x_2 + 12x_3 + 9x_4 = 15$$

$$-2L_2 : -4x_1 - 6x_2 - 2x_3 + 4x_4 = -2$$

$$4L_3 : 4x_1 + 8x_2 - 20x_3 + 16x_4 = 12$$

$$(\text{Sum})L : 3x_1 + 5x_2 - 10x_3 + 29x_4 = 25$$

- L is a linear combination of L_1 , L_2 , and L_3
- Is $u = (-8, 6, 1, 1)$ a solution of the system?
- Is $u = (-8, 6, 1, 1)$ a solution of the linear combination?

What can you conclude?

Equivalent systems

Theorem

Given two systems of linear equations, say L_1 and L_2 . They have the same solutions iff each equation in L_1 is a linear combination of the equations in L_2 .

Definition

Two systems of linear equations are **equivalent** if they have the same solutions.

Elementary operations

Given a system of linear equations L_1, L_2, \dots, L_m . The following operations are called **elementary operations**.

- **[E1]** Interchange two of the equations

Interchange L_i and L_j or $L_i \leftrightarrow L_j$

- **[E2]** Replace an equation by a nonzero multiple of itself.

Replace L_i by kL_i or $kL_i \rightarrow L_i$

- **[E3]** Replace an equation by the sum of a multiple of another equation and itself.

Replace L_j by $kL_i + L_j$ or $kL_i + L_j \rightarrow L_j$

Theorem

Given a system \mathcal{L} . Let \mathcal{M} be the system obtained from \mathcal{L} by a finite sequence of elementary operations.

Then \mathcal{M} and \mathcal{L} have the same solutions.

Note: Sometimes E_2 and E_3 can be applied in one step:

[E] Replace equation L_j by $kL_i + k'L_j$ (where $k, k' \neq 0$)

$$kL_i + k'L_j \rightarrow L_j$$

How to find a solution of a linear equations system?

- Use elementary operations to transform the given system into an equivalent system whose solution can be easily obtained

This is called **Gaussian Elimination** (will be discussed later).

Part 4: Small square systems of linear equations

Linear equation in **one variable**

Example

Solve the following linear system of one variable:

- $4x - 1 = x + 6$
- $2x - 5 - x = x + 3$
- $4 + x - 3 = 2x + 1 - x$

What can you conclude?

Linear equation in **one variable**

Example

Solve the following linear system of one variable:

- $4x - 1 = x + 6$
- $2x - 5 - x = x + 3$
- $4 + x - 3 = 2x + 1 - x$

What can you conclude?

Theorem

Given the system of unique linear equation $ax = b$.

- 1. If $a \neq 0$, then $x = \frac{b}{a}$ is a unique solution of the system.*
- 2. If $a = 0$, but $b \neq 0$, then the system has no solution.*
- 3. If $a = 0$ and $b = 0$, then every scalar k is a solution of $ax = b$.*

Example

Example

Solve the following linear system of one variable:

- $4x - 1 = x + 6$ (Theorem 7 (1))
In standard form: $3x = 7$. Then $x = \frac{7}{3}$ is the unique solution.
- $2x - 5 - x = x + 3$ (Theorem 7 (2))
In standard form: $0x = 8$. The equation has no solution.
- $4 + x - 3 = 2x + 1 - x$ (Theorem 7 (3))
In standard form: $0x = 0$. Then every scalar k is a solution.

System of two linear equations in **two variables**

Given a system of two non-degenerate linear equations in two variables:

$$A_1x + B_1y = C_1$$

$$A_2x + B_2y = C_2$$

Example

Solve the following system of linear equations:

$$\begin{cases} L_1 : x - y = -4 \\ L_2 : 3x + 2y = 12 \end{cases} \quad \begin{cases} L_1 : x + 3y = 3 \\ L_2 : 2x + 6y = -8 \end{cases} \quad \begin{cases} L_1 : x + 2y = 4 \\ L_2 : 2x + 4y = 8 \end{cases}$$

What can you conclude?

The number of solutions of (2×2) -system

1. The system has exactly **one solution**.

$$L_1 : x - y = -4$$

$$L_2 : 3x + 2y = 12$$

2. The system has **no solution**.

$$L_1 : x + 3y = 3$$


$$L_2 : 2x + 6y = -8$$

3. The system has **an infinite number of solutions**.


$$L_1 : x + 2y = 4$$

$$L_2 : 2x + 4y = 8$$


Geometric interpretation



(a) Exactly one solution



(b) No solution



(c) Infinitely many solution

1. System with exactly one solution


- Given:

$$A_1x + B_1y = C_1$$

$$A_2x + B_2y = C_2$$

- Both lines have distinct slopes

$$\frac{A_1}{A_2} \neq \frac{B_1}{B_2} \quad \text{or} \quad A_1B_2 - A_2B_1 \neq 0$$



2. System with no solution


- Given:

$$A_1x + B_1y = C_1$$

$$A_2x + B_2y = C_2$$

- Both lines are parallel (have the same slope)

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} \neq \frac{C_1}{C_2} \quad \text{here } A_1B_2 - A_2B_1 = 0$$



3. System with infinitely many solutions


- Given:

$$A_1x + B_1y = C_1$$

$$A_2x + B_2y = C_2$$

- Both lines have the same slopes and same y-intercepts

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} = \frac{C_1}{C_2} \quad \text{here } A_1B_2 - A_2B_1 = 0$$



Recap

- The system has exactly one solution when $A_1B_2 - A_2B_1 \neq 0$
- The system has no solution of infinitely many solutions when $A_1B_2 - A_2B_1 = 0$

The value $A_1B_2 - A_2B_1$ is called **determinant of order two**

$$\begin{vmatrix} A_1 & B_1 \\ A_2 & B_2 \end{vmatrix}$$

Q: Can you relate the solution of system of linear equations to determinant?

Recap

- The system has exactly one solution when $A_1B_2 - A_2B_1 \neq 0$
- The system has no solution of infinitely many solutions when $A_1B_2 - A_2B_1 = 0$


The value $A_1B_2 - A_2B_1$ is called **determinant of order two**

$$\begin{vmatrix} A_1 & B_1 \\ A_2 & B_2 \end{vmatrix}$$


Q: Can you relate the solution of system of linear equations to determinant?

Remark: A system has a *unique solution* iff the determinant of its coefficients is not zero.

The number of solutions of (3×3) -system



(a) One solution
(a point)




(b) Infinite number
of solutions (a line)



(c) Infinite number
of solutions (a plane)



(d) No solution



(e) No solution

Example 1: unique solution

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 2 & 3 & 1 & 1 \\ 3 & 1 & 2 & 1 \end{array} \right] \xrightarrow{\text{Gaussian elimination}} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 1 \end{array} \right]$$

from which we can derive the set of solution:

$$x_1 = 1, x_2 = 0, x_3 = -1$$

Example 2: infinitely many solution

$$\left[\begin{array}{ccc|c} 1 & 1 & 2 & 4 \\ 2 & -1 & 1 & 2 \\ 1 & 2 & 3 & 6 \end{array} \right] \xrightarrow{\text{Gaussian elimination}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 4 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

From the last row, we can derive the equation:

$$0x_1 + 0x_2 + 0x_3 = 0$$

which can be satisfied by many value of x . The solution can be written in parametric form:

- Let $x_3 = k$, with $k \in \mathbb{R}$
- Then $x_2 = 2 - k$ and $x_1 = 4 - x_2 - 2x_3 = 4 - (2 - k) - 2k = 2 - k$

This means that there are an infinitely many solutions, because there are infinitely many possible values of k .

Example 3: no solution

$$\left[\begin{array}{ccc|c} 1 & 1 & 2 & 4 \\ 2 & -1 & 1 & 2 \\ 1 & 2 & 3 & 7 \end{array} \right] \xrightarrow{\text{Gaussian elimination}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 4 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

From the last row, we can derive the equation:

$$0x_1 + 0x_2 + 0x_3 = 1 \quad (1)$$

Clearly, no possible value of $x_1, x_2, x_3 \in \mathbb{R}$ that can satisfy equation (1).

What about a system with more than 3 variables?

Remark

- For a linear system with more than 3 variables, it's hard to interpret it geometrically.
- However we can check the possible number of solutions by looking at **the shape of the reduced echelon form**.




Figure: Left (unique solution), middle (many solutions), right (no solution) — *source: lecture notes of Rinaldi Munir, ITB*

to be continued...