TUTORIAL VI

1 Binary Erasure Channel

A binary erasure channel with input alphabet $\{0,1\}$ and output alphabet $\{0,1,E\}$ is defined by the following transition probabilities.

$$p_{Y|X}(0|0) = p_{Y|X}(1|1) = 1 - \alpha, \qquad p_{Y|X}(E|0) = p_{Y|X}(E|1) = \alpha$$

Essentially, a fraction α of the input bits are erased (represented by the symbol E).

- 1. Determine the capacity of the channel.
- 2. If there is (noiseless) feedback on whether the input bit is received or erased, how do you achieve a rate equal to the capacity (you can send the same message several times)?
- 3. Suppose that there is no feedback and we use the following coding scheme: encode 0 as 000 and 1 as 111. Decode 000, E00, 0E0, 00E, EE0, E0E, 0EE to 0 and similarly decode 111, E11, 1E1, 11E, EE1, E1E, 1EE to 1. In case EEE is received, then choose one of 0, 1 at random. What is the probability of error for the code?

2 Expurgation

Let C be a M-code with error probability $P_{err}(C) = \delta$.

1. Show that you can build a $\lfloor M/2 \rfloor$ -code with maximal error probability $\leq 2\delta$.

3 Fun with Fano

- 1. Consider the two following pairs of correlated random variables:
 - i. X is uniform on $\{0,1\}^n$, Y equals the first n/2 bits of X.
 - ii. With probability $\alpha \in [0; 1]$, X is uniform on $\{0; 1\}^n$ and Y = X; and with probability 1α , X is uniform on $\{0; 1\}^n$ and Y is the all 0s string.

Suppose we observe Y and estimate $\hat{X} = g(Y)$. What is the minimum possible value of $\mathbf{P}(\hat{X} \neq X)$ in the above two examples ? What lower bound does Fano's inequality give in the two examples ?

2. For two vectors $u, v \in \{0, 1\}^n$, we denote by $\Delta(u, v)$ the following set: $\Delta(u, v) = \text{Card}(\{j \in \{1, \dots, n\} : u_j \neq v_j\})$. Suppose X and Y are two correlated random variables taking values in $\{0, 1\}^n$.

For $i \in \{0, ..., n\}$, we define $\theta_i = \mathbf{P}(\Delta(X, Y) = i)$. Prove that

$$H(X|Y) \le \sum_{i=0}^{n} \theta_i \log_2 \left(\binom{n}{i} \frac{1}{\theta_i} \right)$$

(Hint: Define the random variable $\Delta(X,Y)$ and mimic steps from the proof of Fano's inequality)